

APPM 1345

Exam 2

Spring 2024

Name		
Instructor	Richard McNamara	Section 150

This exam is worth 100 points and has **4 problems**.

**Make sure all of your work is written in the blank spaces provided.** If your solutions do not fit, there is additional space at the end of the test. Be sure to **make a note** indicating the page number where the work is continued or it will **not** be graded.

**Show all work and simplify your 5ua003 work is continued or it**

1. (25 pts) Parts (a) and (b) are unrelated.

- (a) Find the average value  $f_{\text{ave}}$  of the function  $f(x) = 9 - x^2$  on the interval  $[0; 3]$ , and find all values of  $c$  on  $[0; 3]$  for which  $f(c) = f_{\text{ave}}$ .

**Solution:**

$$\begin{aligned} f_{\text{ave}} &= \frac{1}{3} \int_0^3 (9 - x^2) dx = \frac{1}{3} \left[ 9x - \frac{x^3}{3} \right]_0^3 \\ &= \frac{1}{3} (27 - 9) = \boxed{6} \end{aligned}$$

*f* (on the interval  $[0; 3]$ )

2.



4. (28 pts) Parts (a) and (b) are unrelated.

(a) Consider the function  $h(x) = \cos^2 x$  on the interval  $I = [0; -2]$ .

- i. Determine the numerical value of the Riemann sum  $L_2$  for  $h(x)$  on  $I$  using **left** endpoints and 2 equally-sized subintervals. Fully simplify your answer.
- ii. Write an expression for the general Riemann sum  $L_n$  for  $h(x)$  on  $I$  using **left** endpoints and  $n$  equally-sized subintervals. Express your answer using sigma notation.

**Solution:**

$$i. \quad \Delta x = \frac{b - a}{n} = \frac{-2 - 0}{2} = -\frac{1}{2}$$

Since left endpoints are being used and the subintervals are of equal size, we have  $x_0 = 0$  and  $x_1 = -\frac{1}{2}$ .

$$\begin{aligned} L_2 &= [h(x_0) + h(x_1)] \Delta x \\ &= [\cos^2(0) + \cos^2(-\frac{1}{2})] \left(-\frac{1}{2}\right) \\ &= [1 + \frac{3}{4}] \left(-\frac{1}{2}\right) \\ &= \frac{7}{4} \left(-\frac{1}{2}\right) = -\frac{7}{8} \end{aligned}$$

$$ii. \quad \Delta x = \frac{b - a}{n} = \frac{-2 - 0}{n} = -\frac{2}{n}$$

Since left endpoints are being used and the subintervals are of equal size, we have

$$x_{i-1} = a + (i-1) \Delta x = 0 + (i-1) \left(-\frac{2}{n}\right) = -\frac{2(i-1)}{n}$$

$$\begin{aligned} L_n &= \sum_{i=1}^n h(x_{i-1}) \Delta x \\ &= \sum_{i=1}^n \cos^2(x_{i-1}) \left(-\frac{2}{n}\right) \\ &= -\frac{2}{n} \sum_{i=1}^n \cos^2\left(-\frac{2(i-1)}{n}\right) \end{aligned}$$

(b) Suppose the following expression is a Riemann sum for a continuous function  $u(x)$  on the interval  $[1; 2]$ :

$$R_n = \sum_{i=1}^n \left( \frac{3i^2}{n} + 1 \right) \frac{3}{n}$$

Find the numerical value of  $\int_1^2 u(x) dx$  by evaluating the appropriate limit of  $R_n$ . Do not use a Dominance