APPM 1350

Exam 3

Fall 2023

Name	
Instructor	Lecture Section

1.

2. (20 pts) The following problems are unrelated. For each, be sure to fully simplify your answers.

   (a) An object moves along an axis starting at time $t = 0$. It moves with acceleration $\sin(t)$ meters per second squared with an initial velocity of 3 meters per second and an initial position of 2 meters. Find the position of this object at $t = \frac{3\pi}{2}$ seconds. (Provide an exact answer in terms of π. Do not round or approximate your final answer.)

   (b) Consider the integral $\int_0^2 x^3\,dx$. In this problem, you will evaluate this integral in two different ways.

      i. Evaluate this integral using the Fundamental Theorem of Calculus Part 2 (also known as the Evaluation Theorem).

      ii. Separately, write the integral as the limit of a right-hand Riemann sum using n rectangles of equal width. Evaluate this sum and limit without using your work from (i).

3. (10 pts) The following is a graph of $y = f(x)$. Be sure to scan and submit this graph as part of your answer because you will draw on it in addition to answering a few questions (below).

(a) What is the x-value of the only zero of f? (Note: The only zero of f is shown in the above graph.)

(b) With $x_1 = -2$, after one iteration of Newton's Method will x_2 be closer to or farther away from the zero in part (a)?

i. Answer 'closer' or 'farther.'

ii. On the provided graph draw the relevant line supporting your answer from (b). *Label this line as (b).*

(c) Give an x-value x_1 for which Newton's method definitely will not converge.

(d) With $x_1 = 3$, draw a line on the provided graph illustrating one step of Newton's Method. *Label this line as (d).*

(e) What will $\lim_{n \to \infty} x_n$ equal when $x_1 = 3$? (That is, with an initial guess of $x_1 = 3$ and repeated application of Newton's Method, to what value will the x_n's approach?)

3. (10 pts) The following is a graph of $y = f(x)$. Be sure to scan and submit this graph as part of your answer because you will draw on it in addition to answering a few questions (below).

(a) What is the x-value of the only zero of f? (Note: The only zero of f is shown in the above graph.)

(b) With $x_1 = -2$, after one iteration of Newton's Method will x_2 be closer to or farther away from the zero in part (a)?

i. Answer `closer' or `farther.'

ii. On the provided graph draw the relevant line supporting your answer from (b). Label this line as (b).

(c) Give an x-value x_1 for which Newton's method definitely will not converge.

(d) With $x_1 = 3$, draw a line on the provided graph illustrating one step of Newton's Method. Label this line as (d).

(e) What will $\lim_{n \to \infty} x_n$ equal when $x_1 = 3$? (That is, with an initial guess of $x_1 = 3$ and repeated application of Newton's Method, to what value will the x_n's approach?)

2. (20 pts) The following problems are unrelated. For each, be sure to fully simplify your answers.

(a) An object moves along an axis starting at time $t = 0$. It moves with acceleration $\sin(t)$ meters per second squared with an initial velocity of 3 meters per second and an initial position of 2 meters. Find the position of this object at $t = \frac{3}{2}$ seconds. (Provide an exact answer in terms of π. Do not round or approximate your final answer.)

(b) Consider the integral $\int_0^2 x^3 \, dx$. In this problem, you will evaluate this integral in two different ways.

i. Evaluate this integral using the Fundamental Theorem of Calculus Part 2 (also known as the Evaluation Theorem).

ii. Separately, write the integral as the limit of a right-hand Riemann sum using n rectangles of equal width. Evaluate this sum and limit without using your work from (i).

2. (20 pts) The following problems are unrelated. For each, be sure to fully simplify your answers.

(a) An object moves along an axis starting at time $t = 0$. It moves with acceleration $a(t) = 2t$ meters per second squared with an initial velocity of 3 meters per second and an initial position of 2 meters. Find the position of this object at $t = \frac{1}{2}$ seconds. (Provide an exact answer in terms of π. Do not round or approximate your real answer.)

(b) Consider the integral $\int_0^2 x^3 \, dx$. In this problem, you will evaluate this integral in two different ways.

i. Evaluate this integral using the Fundamental Theorem of Calculus Part 2 (also known as the Evaluation Theorem).

ii. Alternately, write the integral as the limit of a right-hand Riemann sum using n rectangles of equal width. Evaluate this sum and limit without using your work from (i).

2. (20 pts) The following problems are unrelated. For each, be sure to fully simplify your answers.

   (a) An object moves along an axis starting at time $t = 0$. It moves with acceleration $a(t)$ meters per second per second and initial velocity of 3 meters per second and an initial position of 2 meters. Find the position of this object at $t = \frac{\pi}{2}$ seconds. (Provide an exact answer in terms of π do not round or approximate your final answer.)

   (b) Consider the integral $\int_0^2 x^3\,dx$. In this problem, you will evaluate this integral in two different ways.

      i. Evaluate this integral using the Fundamental Theorem of Calculus Part 2 (also known as the Evaluation Theorem).

      ii. Separately, write the integral as the limit of a right-hand Riemann sum using n rectangles of equal width. Evaluate this sum and limit without using your work from (i).